arXiv:0901.3588v2 [hep-th] 4 Apr 2009

Quasinormal modes of warped AdS_3 black holes and warped AdS/CFT correspondence

Bin Chen and Zhi-bo Xu*

Department of Physics

and State Key Laboratory of Nuclear Physics and Technology,

Peking University, Beijing 100871, P.R.China

Abstract

We calculate the scalar quasinormal modes of warped AdS_3 black holes analytically. We find that in general they are not in agreement with the prediction of the usual AdS/CFT correspondence. Nevertheless, for the black hole not deviating far from BTZ black hole, the quasinormal frequencies are still well consistent with AdS/CFT correspondence.

---

*Email:bchen01,xuzhibo@pku.edu.cn

The quasinormal modes of black holes have been studied in general relativity for nearly forty years (please see [1] for excellent reviews and detailed list of references). They characterize the "sound" of the black holes. For the perturbations of the black holes, they obey the linearized equations of motion. And the quasinormal modes are defined as the perturbations subject to the physical boundary conditions that near the horizon of the black holes the local solution is purely ingoing and at spatial infinity the solution is purely outgoing. This choice of boundary condition makes generically the frequency of the perturbation be complex. Therefore the quasinormal perturbations undergo damped oscillations, just as the ring of a bell. Physically, the perturbations can fall into the black holes and decay from the point of view of an observer at the infinity. This is true not only for the black holes in flat spacetime and also in anti-de Sitter(AdS) spacetime.

On the other hand, the quasinormal modes of AdS black holes are of particular interests in the study of AdS/CFT correspondence[2]. From AdS/CFT correspondence, the presence of black holes in anti-de Sitter spacetime corresponds to turning on a temperature in dual conformal field theory. The quasinormal modes of black holes correspond to the tiny deviations of the thermal equilibrium in dual field theory, which have nonzero damping[3]. It turns out that the complex frequencies of quasinormal modes can be understood as the poles in the retarded green function in the dual finite temperature field theory[4, 5]. Especially for two-dimensional conformal field theory, the left and right sectors are independent. At thermal equilibrium, the two sectors may have different temperatures (T_L, T_R). Consider a small perturbation operator $\mathcal{O}$ with conformal weights (h_L, h_R). Under such a perturbation, the system will return to thermal equilibrium exponentially with a characteristic time scale, which is inversely proportional to the imaginary part of the poles of the correlation function of the operator $\mathcal{O}$ in momentum space. There are two sets of poles

$$\begin{aligned}\omega_L &= k - 4\pi i T_L(n + h_L) \\ \omega_R &= -k - 4\pi i T_R(n + h_R), \qquad (1)\end{aligned}$$

where n being non-negative integer. In [4], it has been shown that these poles are in exact agreement with the quasinormal frequencies of the BTZ black hole.

The Banados-Teitelboim-Zanelli(BTZ) black hole is a solution of the vacuum Einstein equations in three-dimensional anti-de Sitter spacetime[6]. The metric of the BTZ black hole is of the form

$$\begin{aligned}ds^2 &= -(-M + \frac{\rho^2}{l^2} + \frac{J^2}{4\rho^2})d\tau^2 + (-M + \frac{\rho^2}{l^2} + \frac{J^2}{4\rho^2})^{-1}d\rho^2 \\ &\quad + \rho^2(d\phi - \frac{J}{2\rho^2}d\tau)^2, \qquad (2)\end{aligned}$$

in Schwarzschild coordinates, where l is the radius of AdS_3 spacetime and M, J are the mass and angular momentum of the black hole. There are two horizons in BTZ black hole, outer one ρ_+ and inner one ρ_- decided by M, J. BTZ black holes are asymptotic to AdS_3 spacetime. According to AdS_3/CFT_2 dictionary, the two independent temperatures are related to the horizons

$$T_L = \frac{\rho_+ - \rho_-}{2\pi l}, \qquad T_R = \frac{\rho_+ + \rho_-}{2\pi l}. \tag{3}$$

For a scalar field with mass m in AdS_3, its corresponding operator in CFT_2 has conformal weights

$$h_L = h_R = \frac{1}{2}(1 + \sqrt{1 + m^2 l^2}). \tag{4}$$

The quasinormal modes of the BTZ black hole were computed for the first time in [7]. And it was found in [4] that the scalar quasinormal frequencies of BTZ black hole were in precise match with (1).

The warped AdS_3 black hole is the vacuum solution in three dimensional topologically massive gravity (TMG) with a negative cosmological constant. It was first discovered in [8]. Very recently, it was found that just as BTZ black holes are discrete quotients of AdS_3 spacetime, the warped AdS_3 black holes are discrete quotients of warped AdS_3 spacetime[9]. It was even conjectured in [9] that the warped black holes are holographically dual to a two-dimensional conformal field theory. The study of thermodynamics of the warped black holes gave strong support of the conjecture. It would be interesting to investigate this nontrivial AdS_3/CFT_2 correspondence further[1]. For other recent works on warped AdS_3 black holes, see [10].

In this letter, we study the scalar quasinormal modes of the warped AdS_3 black holes and check if it is consistent with the prediction of AdS/CFT correspondence.

Let us start from the spacelike stretched black holes, discussed in [8, 9]. Its metric takes the following form in terms of Schwarzschild coordinates:

$$ds^2 = -N(r)^2 dt^2 + R(r)^2 [d\theta + N^\theta(r) dt]^2 + \frac{l^2 dr^2}{4R(r)^2 N(r)^2}, \tag{5}$$

where

$$\begin{aligned} R(r)^2 &= \frac{r}{4}\left(3(v^2-1)r + (v^2+3)(r_+ + r_-) - 4v\sqrt{r_+ r_- (v^2+3)}\right) && (6) \\ N^2(r) &= \frac{(v^2+3)(r-r_+)(r-r_-)}{4R(r)^2}, && (7) \\ N^\theta(r) &= \frac{2vr - \sqrt{r_+ r_- (v^2+3)}}{2R(r)^2}, && (8) \end{aligned}$$

---

[1] It would be better to call it as warped AdS_3/CFT_2 correspondence, since the conformal boundary of warped AdS_3 is different from the one of AdS_3. We thank the referee for this suggestion.

where $-l^{-2}$ is the negative cosmological constant and the parameter $v = \mu l/3$ with μ being the mass of the graviton. Just like the BTZ black hole, there are two horizons located at $r = r_+$ and $r = r_-$. We will focus on the physical black holes, in which case $v \geq 1$. When $v = 1$, there is no stretching and the above black hole becomes the usual BTZ black hole. In [9], the temperatures of the black holes were identified to be

$$T_H = \frac{4\pi v l}{v^2+3}\frac{T_L+T_R}{T_R}, \tag{9}$$

where

$$\begin{aligned} T_L &= \frac{(v^2+3)}{8\pi l}\left(r_+ + r_- - \frac{\sqrt{(v^2+3)r_+r_-}}{v}\right) && (10) \\ T_R &= \frac{(v^2+3)(r_+-r_-)}{8\pi l} && (11) \end{aligned}$$

are the temperature of dual CFT. The dual two-dimensional CFT is supposed to have the central charges

$$c_L = \frac{l}{G}\frac{4v}{v^2+3}, \qquad c_R = \frac{l}{G}\frac{5v^2+3}{v(v^2+3)}. \tag{12}$$

When $v = 1$, the warped black holes is the BTZ black holes. This could be seen by a coordinate transformation

$$t = \frac{\rho_+ - \rho_-}{l}\tau, \quad \theta = \phi - \frac{1}{l}\tau, \quad r = \frac{\rho^2}{\rho_+ - \rho_-}. \tag{13}$$

In terms of the coordinates r, the horizons are

$$\rho_\pm = \sqrt{r_\pm}(\sqrt{r_+} - \sqrt{r_-}). \tag{14}$$

Note that the definition of the temperatures is actually related to the coordinate choice. In BTZ case, one has

$$\begin{aligned} T_L &= \frac{\rho_+ - \rho_-}{2\pi l} = \frac{(\sqrt{r_+} - \sqrt{r_-})^2}{2\pi l} \\ T_R &= \frac{\rho_+ + \rho_-}{2\pi l} = \frac{r_+ - r_-}{2\pi l}, \end{aligned}$$

which is the same as (10). However, notice that the transformation (13) relate two time direction in two different coordinates so that the temperatures in two coordinates should be related by

$$T(r) = \frac{l}{\rho_+ - \rho_-}T(\rho). \tag{15}$$

Taking into account of this subtlety, one has

$$\begin{aligned} \tilde{T}_L &= \frac{1}{2\pi}, \\ \tilde{T}_R &= \frac{1}{2\pi}\frac{\sqrt{r_+}+\sqrt{r_-}}{\sqrt{r_+}-\sqrt{r_-}}. \end{aligned} \tag{16}$$

To study the quasinormal modes of the blackhole, we consider the scalar perturbation in the background for simplicity. The scalar wavefunction satisfies the equation

$$(\nabla_\mu \nabla^\mu - m^2)\Phi = 0 \tag{17}$$

Since the background has the translational isometry along t and θ, we may make the following ansatz

$$\Phi = e^{-i\omega t + ik\theta}\phi, \tag{18}$$

to simplify the equation, and we have

$$\frac{4}{l^2}R^2N^2\partial_r(R^2N^2\partial_r)\phi + (R^2\omega^2 + 2k(R^2N^\theta)\omega + k^2 - m^2N^2R^2)\phi = 0. \tag{19}$$

Define the variable

$$z = \frac{r-r_+}{r-r_-}, \tag{20}$$

and we are led to the following equation after a lengthy calculation

$$z(1-z)\frac{d^2\phi}{dz^2} + (1-z)\frac{d\phi}{dz} + \frac{1}{(v^2+3)^2}\left(\frac{A}{z} + B + \frac{C}{1-z}\right)\phi = 0, \tag{21}$$

where

$$A = \frac{l^2}{(r_+-r_-)^2}(2k + \omega\sqrt{r_+}(2v\sqrt{r_+} - \sqrt{v^2+3}\sqrt{r_-}))^2, \tag{22}$$

$$B = -\frac{l^2}{(r_+-r_-)^2}(2k + \omega\sqrt{r_-}(2v\sqrt{r_-} - \sqrt{v^2+3}\sqrt{r_+}))^2, \tag{23}$$

$$C = 3(v^2-1)\omega^2 - m^2(v^2+3). \tag{24}$$

The equation (21) could be transformed to standard hypergeometric function form. Near the horizon $z = 0$ or $r = r_+$, there are two independent solutions

$$\phi_1 = z^\alpha(1-z)^\beta F(a,b,c,z), \qquad \phi_2 = z^\alpha(1-z)^\beta F(a+c-1, b-c+1, 2-c, z) \tag{25}$$

where

$$\begin{aligned} \alpha &= -i\frac{\sqrt{A}}{v^2+3}, \\ \beta &= \frac{1}{2}\left(1 - \sqrt{1 - \frac{4Cl^2}{(v^2+3)^2}}\right), \end{aligned}$$

and

$$
\begin{aligned}
c &= 2\alpha + 1, \\
a &= \alpha + \beta + i\sqrt{-B}/(v^2+3), \\
b &= \alpha + \beta - i\sqrt{-B}/(v^2+3).
\end{aligned}
$$

Since by definition the quasinormal modes have to be purely ingoing at the horizon, ϕ_1 has the right boundary behavior. And in usual anti-de Sitter spacetime the boundary condition that at the infinity the solution should vanish. For the warped AdS_3 spacetime, the boundary condition at infinity is still under investigation[10, 11]. It is not natural to impose the vanishing Dirichlet boundary condition at radial infinity, since the conformal boundary of warped AdS_3 is very different from the one of usual AdS_3. Actually one may analyze the asymptotic differential equation for radial wavefunction and find that the potential is not infinitely high at radial infinity. Therefore, instead of choosing some kind of boundary condition on wavefunction directly at infinity, we impose the physical requirement that the wavefunction is just purely outgoing at infinity and its corresponding flux is finite. The flux is

$$\mathcal{F} = \sqrt{-g}g^{rr}\frac{1}{2i}(\Phi^*\partial_r\Phi - \Phi\partial_r\Phi^*) \propto (r-r_+)(r-r_-)(\Phi^*\partial_r\Phi - \Phi\partial_r\Phi^*) \tag{26}$$

If ω is a ordinary complex number with both real and imaginary parts, then β is a complex number and not real. We require $\mathrm{Re}\sqrt{1-\frac{4Cl^2}{(v^2+3)^2}} \geq 0$ to avoid any ambiguity. So the asymptotic flux at the infinity has a set of divergent terms and the leading term is of order $(1-z)^{2\beta-1}$, where $\mathrm{Re}2\beta - 1 < 0$. If ω is a purely imaginary number, then β is real and $\beta < 0$ and the leading order divergent term is $(1-z)^{2\beta}$. The leading term are all proportional to

$$\left|\frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}\right|^2 \tag{27}$$

Thus, if we require the asymptotic flux at the infinity is not divergent, we have to set

$$c - a = -n, \qquad \text{or} \qquad c - b = -n, \tag{28}$$

with n being a non-negative integer. Note that these two relations could be obtained by simply imposing vanishing Dirichlet condition at infinity. Let us analyze them case by case.

1) Case 1: $c - a = -n$

In this case, we are led to the following equation on ω:

$$-i\frac{l}{r_+ - r_-}\frac{1}{v^2+3}(4k + \omega\delta) + \frac{1}{2}\left(1 + \sqrt{1 - \frac{4Cl^2}{(v^2+3)^2}}\right) = -n, \tag{29}$$

where

$$\delta \equiv 2v(r_+ + r_-) - 2\sqrt{(v^2+3)r_+r_-}. \tag{30}$$

The solution of (29) is of quite involved form:

$$\omega_R = \frac{v^2+3}{d^2\delta^2 - 3(v^2-1)l^2}\left\{-d\delta\left(\frac{4kd}{v^2+3} + i(n+\frac{1}{2})\right) - i(e+if)\right\}, \tag{31}$$

where

$$d \equiv \frac{l}{r_+ - r_-}. \tag{32}$$

And in the above solution, we have introduced

$$e = \sqrt{\frac{\sqrt{E^2+F^2}+E}{2}}, \qquad f = \sqrt{\frac{\sqrt{E^2+F^2}-E}{2}}, \tag{33}$$

where

$$\begin{aligned} E &= (\frac{1}{4} + \frac{m^2l^2}{v^2+3})d^2\delta^2 - 3(v^2-1)l^2\left((\frac{1}{4}+\frac{m^2l^2}{v^2+3}) + (\frac{4kd}{v^2+3})^2 - (n+\frac{1}{2})^2\right), \\ F &= -3(v^2-1)l^2(n+\frac{1}{2})\frac{8kd}{v^2+3}. \end{aligned}$$

2) Case 2: $c - b = -n$
In this case, the equation on ω is much simpler,

$$-n - \frac{1}{2} + i\frac{2vl\omega}{v^2+3} = \frac{1}{2}\sqrt{1 - \frac{4Cl^2}{(v^2+3)^2}}, \tag{34}$$

which has the solution

$$\omega_L = -i\frac{1}{l}\left\{(2n+1)v + \sqrt{3(n+\frac{1}{2})^2(v^2-1) + (\frac{1}{4}+\frac{m^2l^2}{v^2+3})(v^2+3)}\right\}. \tag{35}$$

Note that the frequency is pure imaginary, being independent of the angular momentum.

The above expression looks quite involved. It would be illuminating to consider the $v = 1$ limit to get a favor and have a consistent check. In the $v = 1$ limit, we have

$$\omega_R = -\frac{2k}{(\sqrt{r_+} - \sqrt{r_-})^2} - i\frac{2(\sqrt{r_+}+\sqrt{r_-})}{l(\sqrt{r_+}-\sqrt{r_-})}\left(n + \frac{1}{2}(1+\sqrt{1+m^2l^2})\right) \tag{36}$$

$$\omega_L = -i\frac{2}{l}\left(n + \frac{1}{2}(1+\sqrt{1+m^2l^2})\right). \tag{37}$$

At first looking, the above result is slightly different from (1) obtained in [4]. The differences reside on the temperature and angular momentum. Both differences originate from the choice of different coordinate systems. For example, taking into account of the coordinate difference, the temperature should be as (16), so that we still have the same physical picture $\omega_{L,R} \propto -\frac{4\pi i}{l}\tilde{T}_{L,R}(1+h_{L,R})$.

The above solution could be simplified in the following limits.

1. In the limit of $v \to \infty$, the expression of the solutions could be simplified to

$$\omega_R = -i\frac{v}{d^2\tilde{\delta}^2 - 3l^2}\left(d\tilde{\delta}(n+\frac{1}{2}) + \sqrt{\frac{1}{4}d^2\tilde{\delta}^2 + 3l^2 n(n+1)}\right), \tag{38}$$

$$\omega_L = -i\frac{v}{l}\left(n+\frac{1}{2}+\sqrt{3(n+\frac{1}{2})^2+1}\right) \tag{39}$$

   where

$$\tilde{\delta} = 2(r_+ + r_- - \sqrt{r_+ r_-}). \tag{40}$$

   The real part of the quasinormal modes and the mass dependence are suppressed in the large v limit. The quasinormal frequencies look pure imaginary and are proportional to v. Physically this means that since the spacetime is stretched so much, the relaxation of the system to thermal equilibrium is extremely fast.

2. For the extremal black hole with $r_+ = r_-$, the fact that $d \to \infty$ leads to

$$\omega_R = -\frac{4k}{\delta}. \tag{41}$$

   Therefore, there is no imaginary part no matter how large v is. This may indicates that the right temperature is zero. And the left temperature could be taken as the constant $\frac{v}{2\pi}$. We note that this phenomenon is similar to the case in Kerr/CFT correspondence[11], where $T_R = 0, T_L = 1/2\pi$.

   Another subtle point arise in the extremal BTZ black hole. From our calculation, it seems that $\tilde{T}_R$ in (16) diverge. However the temperatures in (3) is finite. This indicates that the coordinate transformation is singular and the definition of the temperature is closely related to the choice of the coordinates[12].

Let us return to the general case. The expression of the right frequency looks awful. To simplify the discussion, we may let the angular momentum vanishing $k = 0$. Then we have

$$\begin{aligned}\omega_R = & -i\frac{v^2+3}{d^2\delta^2 - 3(v^2-1)l^2}\Big\{(n+\frac{1}{2})d\delta \\ & +\frac{1}{2}\sqrt{(1+\frac{4m^2l^2}{v^2+3})\left(d^2\delta^2 - 3(v^2-1)l^2\right) + 3(2n+1)^2(v^2-1)l^2}\Big\}.\end{aligned} \tag{42}$$

With the expression of ω_L, it seems that the quasinormal frequencies is not in agreement with the prediction (1) from AdS/CFT correspondence.

However, we have a few remarks:

- One interesting point is that in the above relations, there is a combination
$$1+\frac{4m^2l^2}{v^2+3} \tag{43}$$
in the squared root. This reminds us of the conformal weight of the scalar field with mass m in anti-de Sitter spacetime with radius $2l/\sqrt{v^2+3}$.

- If we can neglect the ω^2 term in C, we have $C=-m^2(v^2+3)$ so that
$$\begin{aligned}\omega_L &= -i\frac{1}{l}\frac{v^2+3}{2v}\left(n+\frac{1}{2}\tilde{\Delta}\right), &(44)\\ \omega_R &= -\frac{4k}{\delta}-i\frac{v^2+3}{l}\frac{r_+-r_-}{\delta}\left(n+\frac{1}{2}\tilde{\Delta}\right), &(45)\end{aligned}$$
with
$$\tilde{\Delta}=1+\sqrt{1+\frac{4m^2l^2}{v^2+3}}. \tag{46}$$
This is actually in exact match with the prediction (1) of AdS_3/CFT_2 correspondence. Firstly note that $\tilde{\Delta}$ is the scaling dimension of the dual operator corresponding to the scalar field of mass m in AdS_3 spacetime with radius $2l/\sqrt{v^2+3}$. Secondly note that for scalar perturbation in CFT, one has $h_L=h_R=\tilde{\Delta}/2$. Thirdly the temperature has to be modified with respect to coordinate transformation. In [9], the temperatures are identified as
$$T_L=\frac{1}{2\pi l}\frac{v^2+3}{8v}\delta, \qquad T_R=\frac{1}{2\pi l}\frac{v^2+3}{4}(r_+-r_-). \tag{47}$$
And the temperatures in the quasinormal frequencies are
$$\tilde{T}_{L,R}=T_{L,R}/\delta. \tag{48}$$
This is what also happens in BTZ case.

  We need to justify our approximation. In order to have $3(v^2-1)\omega^2 << m^2(v^2+3)$, one may expect that in the very low frequency limit $|\omega|<<m$, the above approximation make sense. However, the expression (44) could not be in consistent with this limit. The other possibility is when v is very close to 1. In this case, the black hole solutions are very much like BTZ black holes. In a sense, the BTZ black hole is the only one which can exactly match the AdS/CFT prediction. A slightly deviation from BTZ black hole is still in good agreement with AdS/CFT correspondence.

- There is a singular point at which the right-moving quasinormal mode is not well-defined. Notice that
$$d^2\delta^2-3(v^2-1)l^2=\frac{l^2}{(r_+-r_-)^2}(\sqrt{v^3+3}(r_++r_-)-4v\sqrt{r_+r_-})^2, \tag{49}$$

which vanishes when

$$\frac{r_-}{r_+} = \frac{2v - \sqrt{3(v^2-1)}}{2v + \sqrt{3(v^2-1)}}. \tag{50}$$

This relation is in consistent with the fact that $0 < r_-/r_+ \leq 1$. However, in this case, from (31) the right-moving quasinormal mode has a negative infinite imaginary part, indicating its damping time is zero. The AdS/CFT correspondence suggest that the dual right temperature is infinite, which is not true even after taking into account of the redefinition of the temperature due to coordinate choice. Note that this issue does not arise in the $v = 1$ limit.

- The discrepancy between semi-classical gravity calculation and the prediction of AdS/CFT correspondence may originate from the deformation of AdS_3 spacetime at asymptotic region. The warped AdS_3 black hole is asymptotic to the warped AdS_3 spacetime, in which case the AdS/CFT correspondence has not been well understood. For example, we have no idea of the exact relation between the masses of scalar fields in the bulk and the conformal weight (h_L, h_R) of corresponding operators in CFT. The usual way to decide the relation from supergravity calculation cannot be applied here since the boundary of the warped AdS_3 is not the spacetime in which the CFT_2 is defined. Furthermore, our study suggest that if there does exist a warped AdS/CFT correspondence, the identification of the quasinormal modes of the warped AdS_3 blackhole with the poles in the retarded Green funciton in the dual field theory may not be exact. Obviously a further thorough study of warped AdS/CFT correspondence is deserving.

Acknowledgments

We would like to thank Wei Song for very valuable discussions. The work was partially supported by NSFC Grant No.10535060,10775002 and NKBRPC (No. 2006CB805905).

References

[1] H.P. Nollert, *Quasinormal modes: the characteristic 'sound' of black holes and neutron stars*, Class. Quan. Gravity **16**, R159 (1999).
K.D. Kokkotas and B. G. Schmidt, *Quasi-normal modes of stars and black holes*, Living. Rev. Rel. **2** (1999)2.

[2] J.Maldacena, *The large N limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2** (1998)231, [hep-th/9711200].
S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from non-critical string theory*, Phys. Lett. B **428**, 105 (1998), [hep-th/9802109].
E. Witten, *Anti-de Sitter space and holography*, Adv. Theor. Math. Phys. **2**, 253 (1998), hep-th/9802150.

[3] G.T. Horowitz and V.E. Hubeny, *Quasinormal modes of AdS black holes and the approach to thermal equilibrium*, Phys. Rev. D **62**, 024027 (2000), [hep-th/9909056].

[4] D. Birmingham, *Choptuik scaling and quasinormal modes in the anti-de Sitter space/conformal field theory correspondence*, Phy. Rev. D**64**, 064024 (2001).
D. Birmingham, I. Sachs and S.N. Solodukhin, *Conformal field theory interpretation of black hole quasinormal modes*, Phys. Rev. Lett. **88**, 151301 (2002), [hep-th/0112055].

[5] D.T. Son and A.O. Stariets, *Minkowski-space correlators in AdS/CFT correspondence: Recipe and applications*, JHEP **0209**, 042 (2002), [hep-th/0205051].

[6] M. Banados, C. Teitelboim and J. Zanelli, *The black hole in three-dimensional space-time*, Phys. Rev. Lett. **69**, 1849 (1992), [hep-th/9204099].

[7] V. Cardoso and J. P. S. Lemos, *Scalar, electromagnetic and Weyl perturbations of BTZ black holes: Quasinormal modes*, Phys. Rev.D**63**, 124015 (2001) [arXiv:gr-qc/0101052].

[8] A. Bouchareb and G. Clement, *Black hole mass and angular momentum in topologically massive gravity*, Class. Quant. Grav. **24**, 5581 (2007), [arXiv:0706.0263].

[9] D. Anninos, W. Li, M. Padi, W. Song and A. Strominger, *Warped AdS_3 black holes*, [arXiv:0907.3040].

[10] G. Compere and S. Detournay, *Semi-classical central charge in topologically massive gravity*, [arXiv:0808.1911].G. Compere, S. Detournay and M. Romo, *Supersymmetric Godel and warped black holes in string theory*, Phys. Rev. D**78**:104030,2008, [arXiv:0808.1912]. D. Anninos, *Hopfing and Puffing Warped Anti-de Sitter Space*, [arXiv:0809.2433]. J. Oh and W. Kim, *Absorption Cross Section in Warped AdS_3 Black Hole*, [arXiv:0811.2632].

[11] M. Guica, T. Hartman, W. Song and A. Strominger, *The Kerr/CFT correspondence*, [arXiv:0809.4266].

[12] W. Kim and E.J. Son, *Thermodynamics of warped* AdS_3 *black hole in the brick wall method*, [arXiv: 0812.0876].